

INTEGRAL COHOMOLOGY AND MIRROR SYMMETRY FOR CALABI-YAU 3-FOLDS

VICTOR BATYREV AND MAXIMILIAN KREUZER

ABSTRACT. In this paper we compute the integral cohomology groups for all examples of Calabi-Yau 3-folds obtained from hypersurfaces in 4-dimensional Gorenstein toric Fano varieties. Among 473 800 776 families of Calabi-Yau 3-folds X corresponding to 4-dimensional reflexive polytopes there exist exactly 32 families having non-trivial torsion in $H^*(X, \mathbb{Z})$. We came to an interesting observation that the torsion subgroups in H^2 and H^3 are exchanged by the mirror symmetry involution.

INTRODUCTION

Let X be a smooth projective Calabi-Yau d -fold over $\mathbb{C}$. We shall always assume that $h^i(X, \mathcal{O}_X) = 0$ for $0 < i < d$. Our main interest is the torsion in the integral cohomology ring $H^*(X, \mathbb{Z})$. For a finite abelian group G we denote by G^* the dual abelian group $\text{Hom}(G, \mathbb{Q}/\mathbb{Z})$. Using the isomorphism from the universal coefficient theorem

$$\text{Tors}(H_i(X, \mathbb{Z})) \cong [\text{Tors}(H^{i+1}(X, \mathbb{Z}))]^*$$

and the Poincaré duality

$$\text{Tors}(H_i(X, \mathbb{Z})) \cong \text{Tors}(H^{2d-i}(X, \mathbb{Z})),$$

we obtain for $d = 3$:

$$\begin{aligned} H^0(X, \mathbb{Z}) &\cong H^6(X, \mathbb{Z}) \cong \mathbb{Z}, \quad H^1(X, \mathbb{Z}) = 0, \\ H^2(X, \mathbb{Z}) &\cong A(X) \oplus \mathbb{Z}^a, \quad H^4(X, \mathbb{Z}) \cong B(X)^* \oplus \mathbb{Z}^a \\ H^3(X, \mathbb{Z}) &\cong B(X) \oplus \mathbb{Z}^{2b+2}, \quad H^5(X, \mathbb{Z}) \cong A(X)^*, \end{aligned}$$

where the finite abelian groups

$$A(X) := \text{Tors}(H^2(X, \mathbb{Z})), \quad B(X) := \text{Tors}(H^3(X, \mathbb{Z}))$$

determine completely the torsion in $H^*(X, \mathbb{Z})$. In particular, we have the isomorphisms

$$A(X) \oplus B(X)^* \cong \text{Tors}(H^{\text{even}}(X, \mathbb{Z})) \cong [\text{Tors}(H^{\text{odd}}(X, \mathbb{Z}))]^*.$$

It is known that mirror symmetry exchanges the integers a, b so that for a mirror Calabi-Yau 3-fold X^* one has

$$\begin{aligned} \mathbb{Q}^{2a+2} &\cong H^{\text{even}}(X, \mathbb{Q}) \cong H^{\text{odd}}(X^*, \mathbb{Q}), \\ \mathbb{Q}^{2b+2} &\cong H^{\text{odd}}(X, \mathbb{Q}) \cong H^{\text{even}}(X^*, \mathbb{Q}). \end{aligned}$$

An interesting mathematical question is to understand the behavior of the torsion in $H^*(X, \mathbb{Z})$, i.e. the groups $A(X)$ and $B(X)$, under the mirror symmetry. The group $B(X)$ is isomorphic to the torsion in the étale cohomology group $H_{\text{et}}^2(X, \mathcal{O}_X^*)$ and it is called the *cohomological Brauer group of X* . By a recent result of Kresch and Vistoli [16], $B(X)$ is also isomorphic to the Brauer group of Azumaya algebras on X . On the other hand, there are canonical isomorphisms

$$A(X) \cong \text{Hom}(\pi_1(X), \mathbb{Q}/\mathbb{Z}) \cong \text{Tors}(\text{Pic}(X)).$$

From the physical point of view it is more natural to consider the topological K -groups $K^0(X)$ and $K^1(X)$ together with some natural filtrations [7, 8]. Using the Atiyah-Hirzebruch spectral sequence (see [7], 2.5), one obtains two short exact sequences

$$0 \rightarrow A(X)^* \rightarrow \text{Tors}(K^1(X)) \rightarrow B(X) \rightarrow 0,$$

$$0 \rightarrow B(X)^* \rightarrow \text{Tors}(K^0(X)) \rightarrow A(X) \rightarrow 0.$$

It is expected that the mirror symmetry exchange K^0 and K^1 so that for a mirror pair (X, X^*) one has the isomorphisms

$$\text{Tors}(K^1(X)) \cong \text{Tors}(K^0(X^*)), \quad \text{Tors}(K^0(X)) \cong \text{Tors}(K^1(X^*)).$$

These isomorphisms agree with predictions of SYZ-construction and topological calculations of M. Gross [12]. The compatibility of the above isomorphisms with the natural filtrations in K^i would imply the isomorphisms

$$(1) \quad A(X) \cong B(X^*), \quad B(X) \cong A(X^*).$$

The main purpose of this paper is to verify these isomorphisms for all examples of Calabi-Yau 3-folds obtained from hypersurfaces in 4-dimensional Gorenstein toric Fano varieties.

There exist 473 800 776 families of Calabi-Yau 3-folds X corresponding to 4-dimensional reflexive polytopes [17, 18]. These families give rise to 30 108 different pairs of numbers (a, b) . We show that among 473 800 776 families of Calabi-Yau 3-folds X there exist exactly 32 families having non-trivial torsion in $H^*(X, \mathbb{Z})$. More precisely, there are exactly 16 families of simply-connected Calabi-Yau 3-folds X having non-trivial Brauer group $B(X) \cong \mathbb{Z}/p\mathbb{Z}$ ($p = 2, 3, 5$). They are mirror dual to another 16 families of Calabi-Yau 3-folds X^* having trivial Brauer group and a non-trivial cyclic fundamental group of order $p = 2, 3, 5$ [15]. Thus, we come to the observation that for all families of Calabi-Yau 3-folds one has the isomorphisms (1). Although the groups $A(X)$ and $B(X)$ can be computed in a purely combinatorial way using lattice points in faces of 4-dimensional reflexive polytopes Δ (or dual reflexive polytopes Δ^*) we do not see immediately the isomorphisms (1) from the combinatorial duality between Δ and Δ^* . Therefore, it would

be interesting to find a general mathematical explanation for the mirror symmetry isomorphism

$$B(X) \cong \text{Hom}(\pi_1(X^*), \mathbb{Q}/\mathbb{Z}).$$

If a finite group G acts on a smooth Calabi-Yau manifold V , then there exists a conformal field theory of the orbifold V/G associated with an element $\alpha \in H^2(G, U(1))$ (discrete torsion). In [28] Vafa and Witten have suggested that there should be some connection between elements in the Brauer group of V/G (B -fields) and elements in $H^2(G, U(1))$ (discrete torsion 2-cocycles). In this paper we show that all 16 families of Calabi-Yau 3-folds with non-trivial cyclic Brauer group are birational to Calabi-Yau orbifolds V/G ($G \cong \mathbb{Z}/p\mathbb{Z} \oplus \mathbb{Z}/p\mathbb{Z}$, $p = 2, 3, 5$) and the corresponding elements in the cyclic Brauer group $B(X) \cong H_2(G) \cong \mathbb{Z}/p\mathbb{Z}$ can be identified with the discrete torsion 2-cocycles.

It is easy to see that there are no algebraic curves in X representing the torsion classes in $H_2(X, \mathbb{Z})$. Aspinwall and Morrison suggested in [2] that the instanton expansion of the Yukawa coupling for a Calabi-Yau 3-fold with the Brauer group $B(X) \cong \mathbb{Z}/p\mathbb{Z}$ should be weighted by powers of p -th roots of unity. It would be interesting to verify this prediction for the above 16 toric families of Calabi-Yau 3-folds with non-trivial Brauer group.

Acknowledgements: We would like to thank the Fields Institute for Research in Mathematical Sciences in Toronto for hospitality during the spring of 2005 and DFG-Schwerpunkt “Global methods in complex geometry” for the support of the first author. The second author was supported by FWF grant Nr. P15584-N08. We are grateful to Noriko Yui for her interest and encouragement in our work. We thank also Volker Braun, Mark Gross, Kentaro Hori, Klaus Hulek, Askold Khovanski, Tony Pantev, Duco van Straten, Bernd Siebert and Balázs Szendrői for useful discussions.

1. THE FUNDAMENTAL GROUP OF TORIC HYPERSURFACES

First we recall a combinatorial computation of the fundamental group of a toric variety [9]. Let $M \cong \mathbb{Z}^d$ be a lattice of rank d and $N := \text{Hom}(M, \mathbb{Z})$ the dual lattice. We denote by $\langle *, * \rangle$ the canonical pairing $M \times N \rightarrow \mathbb{Z}$. For an algebraic torus $\mathbb{T}^d := \text{Spec } \mathbb{C}[M] \cong (\mathbb{C}^*)^d$ there are canonical isomorphisms

$$\pi_1^{\text{top}}(\mathbb{T}^d) \cong N \cong H_1(\mathbb{T}^d, \mathbb{Z}), \quad M \cong H^1(\mathbb{T}^d, \mathbb{Z}).$$

If $\mathbb{P}_\Sigma$ is a smooth partial toric compactification of $\mathbb{T}^d$ defined by a fan $\Sigma \subset N_\mathbb{R} = N \otimes \mathbb{R}$, then $\mathbb{P}_\Sigma \setminus \mathbb{T}^d$ is a union of divisors $D_1, \dots, D_n$ which 1-to-1 correspond to primitive lattice vectors $e_1, \dots, e_n \in N$ generating 1-dimensional cones in Σ . If D_i° is a dense torus orbit in D_i , then we have $\mathbb{P}_i := \mathbb{T}^d \cup D_i^\circ \cong \mathbb{C} \times \mathbb{T}^{d-1}$. The embedding $\mathbb{T}^d \hookrightarrow \mathbb{P}_i$ defines the homomorphism of the fundamental groups

$$\rho_i : N = \pi_1^{\text{top}}(\mathbb{T}^d) \rightarrow \pi_1^{\text{top}}(\mathbb{P}_i) \cong N/\mathbb{Z}e_i.$$

Combining all the homomorphisms ρ_i ($i = 1, \dots, n$), we obtain the isomorphisms

$$\pi_1^{\text{top}}(\mathbb{P}_\Sigma) \cong \pi_1^{\text{top}}(\mathbb{P}_{\Sigma^{(1)}}) \cong N / \sum_{i=1}^n \mathbb{Z}e_i,$$

where $\mathbb{P}_{\Sigma^{(1)}}$ is a toric variety associated to the subfan $\Sigma^{(1)} \subset \Sigma$ of all cones of dimension ≤ 1 in Σ . In particular, we have

$$H_1(\mathbb{P}_\Sigma, \mathbb{Z}) \cong H_1(\mathbb{P}_{\Sigma^{(1)}}, \mathbb{Z}) \cong N / \sum_{i=1}^n \mathbb{Z}e_i.$$

In order to compute the fundamental group of hypersurfaces in toric varieties one applies the following result of Oka [22]:

Theorem 1.1. *Let $\Delta \subset M_{\mathbb{R}}$ be a lattice polytope of dimension d ,*

$$f(t) = \sum_{m \in \Delta \cap M} c_m t^m \in \mathbb{C}[M] \cong \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}]$$

be a Δ -nondegenerate Laurent polynomial, and $W \subset \mathbb{T}^d$ the nondegenerate affine hypersurface defined by the equation $f(t) = 0$. Then the embedding $j : W \hookrightarrow \mathbb{T}^d$ is a homotopy $(d-1)$ -equivalence, i.e. the homomorphisms of the homotopy groups

$$j_* : \pi_i(W) \rightarrow \pi_i(\mathbb{T}^d)$$

is bijective for $i < d-1$ and surjective for $i = d-1$.

By Whitehead's theorem, we get:

Corollary 1.2. *The homomorphism*

$$j_* : H_i(W, \mathbb{Z}) \rightarrow H_i(\mathbb{T}^d, \mathbb{Z})$$

is bijective for $i < d-1$ and surjective for $i = d-1$.

Since $H_i(\mathbb{T}^d, \mathbb{Z})$ is a free abelian group of rank $\binom{d}{i}$, using the universal coefficients theorem, we also obtain:

Corollary 1.3. *For any abelian group $\mathcal{A}$, the homomorphism*

$$j^* : H^i(\mathbb{T}^d, \mathcal{A}) \cong \Lambda^i M \otimes_{\mathbb{Z}} \mathcal{A} \rightarrow H^i(W, \mathcal{A})$$

is bijective for $i < d-1$ and injective for $i = d-1$.

Let $\Sigma := \Sigma(\Delta) \subset N_{\mathbb{R}}$ be the normal fan of the polytope $\Delta \subset M_{\mathbb{R}}$. The cones $\sigma = \sigma(\theta) \in \Sigma(\Delta)$ 1-to-1 correspond to faces $\theta \subset \Delta$ in the following way:

$$\sigma(\theta) = \{y \in N_{\mathbb{R}} : \min_{x \in \Delta} \langle x, y \rangle = \langle x, z \rangle \ \forall z \in \theta\},$$

i.e. $\sigma(\theta)$ consists of those $y \in N_{\mathbb{R}}$ for which the minimum of the linear function $\langle *, y \rangle$ on Δ is attained at any point of θ . Sometimes we denote the corresponding projective toric variety $\mathbb{P}_\Sigma = \mathbb{P}_{\Sigma(\Delta)}$ simply by $\mathbb{P}_\Delta$. Let $\overline{W} \subset \mathbb{P}_\Sigma$ be the Zariski closure of W in $\mathbb{P}_\Sigma$. If $\Sigma^{(1)} \subset \Sigma$ is the subfan of all cones of dimension ≤ 1 in Σ , then $\mathbb{P}_{\Sigma^{(1)}}$ and $\overline{W}^{(1)} := \mathbb{P}_{\Sigma^{(1)}} \cap \overline{W}_f$ are smooth quasi-projective varieties.

Theorem 1.4. *Let $d \geq 3$. Then the embedding $j : \overline{W}^{(1)} \hookrightarrow \mathbb{P}_{\Sigma(1)}$ induces the isomorphism*

$$j_* : \pi_1(\overline{W}^{(1)}) \cong \pi_1(\mathbb{P}_{\Sigma(1)}) \cong N / \sum_{i=1}^n \mathbb{Z}e_i,$$

where $e_1, \dots, e_n$ are primitive generators of 1-dimensional cones in $\Sigma^{(1)}$.

Proof. Let $\mathbb{P}_{\Sigma(1)} \setminus \mathbb{T}^d = D_1^\circ \cup \dots \cup D_n^\circ$ ($D_i^\circ \cong \mathbb{T}^{d-1}$). Using the isomorphisms $\pi_1(W) \cong \pi_1(\mathbb{T}^d) \cong N$ and the surjectivity of $\pi_1(W) \rightarrow \pi_1(\overline{W}^{(1)})$, we obtain that

$$j_* : \pi_1(\overline{W}^{(1)}) \rightarrow \pi_1(\mathbb{P}_{\Sigma(1)})$$

is surjective. Therefore the universal covering $\overline{U}^{(1)}$ of $\overline{W}^{(1)}$ is induced by some unramified covering U of $\mathbb{T}^d$.

In order to prove that j_* is an isomorphism it remains to show that the image of every element $e_i \in \pi_1(W) \cong N$ ($i = 1, \dots, n$) is zero in $\pi_1(\overline{W}^{(1)})$. Let $\mathbb{P}_i := \mathbb{T}^d \cup D_i^\circ$ and $\overline{W}_i^{(1)} := \overline{W}^{(1)} \cap \mathbb{P}_i$. Using open inclusions $W \subset \overline{W}_i^{(1)} \subset \overline{W}^{(1)}$, it is enough to prove that the image of $e_i \in \pi_1(W)$ is zero already in $\pi_1(\overline{W}_i^{(1)})$ so that one obtains $\pi_1(\overline{W}_i^{(1)}) \cong N/\mathbb{Z}e_i$. We can consider $\overline{W}_i^{(1)}$ as an affine hypersurface in $\mathbb{P}_i \cong \mathbb{T}^{d-1} \times \mathbb{C} \cong \text{Spec } \mathbb{C}[t_1^{\pm 1}, \dots, t_{d-1}^{\pm 1}, t_d]$ where the divisor $D_i^\circ \subset \mathbb{P}_i$ is defined by $t_d = 0$. Moreover, the element $e_i \in \pi_1(\mathbb{T}^d) = N$ is represented by a small 1-cycle $\gamma_i \subset \mathbb{C}^*$ around $0 \in \mathbb{C}$ which can be contracted to 0 in $\mathbb{T}^{d-1} \times \mathbb{C}$. The transversality of $\overline{W}_i^{(1)}$ and D_i° implies that the 1-cycle γ_i can be chosen inside W_f so that it will be contractible in $\overline{W}_i^{(1)}$. Therefore all elements $e_1, \dots, e_n$ are in the kernel of the homomorphism $\pi_1(W) \rightarrow \pi_1(\overline{W}^{(1)})$. We remark that the isomorphism $\pi_1(\overline{W}_i^{(1)}) \cong \pi_1(\mathbb{P}_i)$ can be interpreted also as a Lefschetz-type statement for a nondegenerate hypersurface $\overline{W}_i^{(1)}$ in $\mathbb{T}^{d-1} \times \mathbb{C}$. $\square$

Since $e_1, \dots, e_n$ generate N over $\mathbb{Q}$, we obtain:

Corollary 1.5. *The fundamental group of the quasi-projective hypersurface $\overline{W}^{(1)}$ is a finite abelian group. In particular, $\pi_1(\overline{W}^{(1)})$ coincides with the algebraic fundamental group of $\overline{W}^{(1)}$ and the universal covering of $\overline{W}^{(1)}$ is a toric quasi-projective hypersurface $\widetilde{W}^{(1)}$ in the finite universal covering of $\mathbb{P}'_{\Sigma(1)} \rightarrow \mathbb{P}_{\Sigma(1)}$ determined by the finite index sublattice $N' = \sum_{i=1}^n \mathbb{Z}e_i \subset N$.*

Theorem 1.6. *Assume that $d \geq 3$. Let $\mathbb{P}_{\widehat{\Sigma}}$ be a smooth d -dimensional projective toric variety defined by a fan $\widehat{\Sigma} \subset N_{\mathbb{R}}$ which is a subdivision of $\Sigma = \Sigma(\Delta)$. We denote by Γ the set of primitive generators v of 1-dimensional cones in $\widehat{\Sigma}^{(1)}$ such that the minimum of $\langle *, v \rangle$ on Δ is attained on a face $\theta \subset \Delta$ of dimension ≥ 1 . Define the sublattice $N_{\Delta}^{(1)} \subset N$ to be generated by all $v \in \Gamma$. If $\widehat{W}$ is the closure of W in $\mathbb{P}_{\widehat{\Sigma}}$, then the fundamental*

group $\pi_1(\widehat{W})$ is isomorphic to the cyclic group

$$N/N_{\Delta}^{(1)}.$$

Proof. Let $\mathcal{E} := \{e_1, \dots, e_n\}$ be the set of primitive generators of 1-dimensional cones in $\Sigma^{(1)}$. Since every linear function $\langle *, e_i \rangle$ attains its minimum of Δ on a $(d-1)$ -dimensional face of Δ , we have $\mathcal{E} \subset \Gamma$ and we can write $\Gamma = \{e_1, \dots, e_n, \dots, e_{n+k}\}$. Moreover, $\overline{W}^{(1)}$ can be considered as a dense open subset of $\widehat{W}$. Therefore, we have a surjective homomorphism

$$\psi : \pi_1(\overline{W}^{(1)}) \rightarrow \pi_1(\widehat{W})$$

and the dual injective homomorphism

$$\psi^* : \text{Hom}(\pi_1(\widehat{W}), \mathbb{Q}/\mathbb{Z}) \rightarrow \text{Hom}(\pi_1(\overline{W}^{(1)}), \mathbb{Q}/\mathbb{Z}).$$

Since $\widehat{W} \setminus \overline{W}^{(1)} = Z_1 \cup \dots \cup Z_k$, where

$$Z_i = D_{n+i} \cap \widehat{W}, \quad i = 1, \dots, k,$$

we obtain that an element $g \in \text{Hom}(\pi_1(\overline{W}^{(1)}), \mathbb{Q}/\mathbb{Z}) \cong \text{Hom}(N/\sum_{i=1}^n \mathbb{Z}e_i, \mathbb{Q}/\mathbb{Z})$ belongs to $\text{Im } \psi^*$ if and only if the cyclic Galois covering of $\overline{W}^{(1)}$ corresponding to g is unramified along all divisors $Z_1, \dots, Z_k$. Using the same arguments as in the proof of 1.4, we obtain that the latter holds exactly when $g(e_{n+1}) = \dots = g(e_{n+k}) = 0$. Therefore,

$$\text{Im } \psi \cong \text{Hom}(N/\sum_{i=1}^{n+k} \mathbb{Z}e_i, \mathbb{Q}/\mathbb{Z}) = \text{Hom}(N/N_{\Delta}^{(1)}, \mathbb{Q}/\mathbb{Z})$$

and by duality $\pi_1(\widehat{W}) \cong N/N_{\Delta}^{(1)}$. The group $N/N_{\Delta}^{(1)}$ is cyclic, because the smoothness of $\widehat{W}$ implies that for any $(d-1)$ -dimensional cone $\sigma = \sigma(\theta) \in \Sigma$ the subset $\sigma \cap \Gamma$ generate a direct summand of the rank $d-1$ of the lattice N (all $(d-1)$ -dimensional cones in $\widehat{\Sigma}$ which subdivide σ are generated by part of a $\mathbb{Z}$ -basis of N). $\square$

Corollary 1.7. *For any lattice point $v \in N$, we denote by*

$$M_v := \{x \in M : \langle x, v \rangle = 0\}$$

the orthogonal complement to v in M . Then we have the exact sequence

$$\bigoplus_{v \in \Gamma} \Lambda^{d-1} M_v \rightarrow \Lambda^{d-1} M \rightarrow H_{\text{tor}}^{2d-3}(\widehat{W}, \mathbb{Z}) \rightarrow 0.$$

Proof. The exactness of

$$\bigoplus_{v \in \Gamma} \Lambda^{d-1} M_v \rightarrow \Lambda^{d-1} M \rightarrow H_{\text{tor}}^{2d-3}(\widehat{W}, \mathbb{Z}) \rightarrow 0$$

follows from the exactness of

$$\bigoplus_{v \in \Gamma} \mathbb{Z}[v] \rightarrow N \rightarrow \pi_1(\widehat{W}) \rightarrow 0$$

together with the canonical isomorphisms

$$\pi_1(\widehat{W}) \cong H_{\text{tor}}^{2d-3}(\widehat{W}, \mathbb{Z}), \quad \Lambda^{d-1}M_v \cong \mathbb{Z}, \quad \Lambda^{d-1}M \cong N$$

after choosing an isomorphism (orientation) $\Lambda^d \cong \mathbb{Z}$. $\square$

Remark 1.8. Theorem 1.6 can be considered as a special case of a more general result of Oka (see [24], Chapter V, §5).

Corollary 1.9. *Let $\overline{W}$ be a nondegenerate $(d-1)$ -dimensional Calabi-Yau hypersurface in the Gorenstein toric Fano variety $\mathbb{P}_\Delta$ associated with a d -dimensional reflexive polytope $\Delta \subset M_{\mathbb{R}}$. Assume $d \geq 3$ and that there exists a smooth projective crepant resolution $\widehat{W}$ of singularities in $\overline{W}$ induced by a convex (coherent) triangulation of the dual reflexive polytope $\Delta^* \subset N_{\mathbb{R}}$. We denote by N'_{Δ^*} the sublattice in N generated by all lattice points in $\Delta^* \cap N$ which belong to faces of codimension > 1 of Δ^* . Then the fundamental group of $\widehat{W}$ is isomorphic to the cyclic group*

$$N/N'_{\Delta^*}.$$

Proof. The statement follows immediately from 1.6, because for a crepant desingularization $\widehat{W} \rightarrow \overline{W}$ the set of primitive generators Γ is exactly the set of all lattice points in faces of codimension < 1 of the dual reflexive polytope Δ^* (see [3]). $\square$

Remark 1.10. Recently Haase and Nill proved that for a reflexive polytope Δ^* of arbitrary dimension $d > 2$ the sublattice in N generated by **all** lattice points in $\Delta^* \cap N$ coincides with N'_{Δ^*} [14]. The proof of this results uses the fact that the interior lattice points in codimension-1 faces of Δ^* are Demazure roots for the automorphism group of the corresponding Gorenstein toric Fano variety $\mathbb{P}_{\Delta^*}$ [20]. In particular, this result easily implies that $N = N'_{\Delta^*}$ for all reflexive polyhedra Δ^* of dimension $d = 3$. The last statement follows also from 1.9 and from the fact that every smooth $K3$ -surface is simply connected.

Remark 1.11. It is known that for $d = 4$ there always exists a smooth projective crepant resolution $\widehat{W}$ of singularities in $\overline{W}$ induced by a convex (coherent) triangulation of the dual reflexive polytope $\Delta^* \subset N_{\mathbb{R}}$. For $d \geq 5$ such a resolution does not exist in general. However, one can use N/N'_{Δ^*} as a candidate for a **stringy fundamental group** of a singular Calabi-Yau variety $\overline{W}$ or its maximal partial crepant desingularization $\widehat{W}$ (similar to stringy Hodge numbers [4]). It would be interesting to know whether for $d \geq 5$ the dual group $(N/N'_{\Delta^*})^*$ coincides with the torsion subgroup in the Picard group of the Deligne-Mumford Calabi-Yau stack associated with the V -manifold $\widehat{W}$ (see [6, 25, 26]).

2. THE LIST OF NON-SIMPLY CONNECTED CALABI-YAU 3-FOLDS

Using the Calabi-Yau data [18] and the program package PALP [19] one can check all 473 800 776 reflexive polytopes Δ and find that there exist exactly 16 examples of reflexive polytopes Δ_i ($1 \leq i \leq 16$) such that $|N/N'_{\Delta_i^*}| > 1$. In the latter case, $|N/N'_{\Delta_i^*}|$ is always a prime number $p = 2, 3, 5$.

The most known example of a non-simply connected Calabi-Yau 3-fold obtained from a hypersurface in a Gorenstein toric variety $\mathbb{P}_\Delta$ is a free μ_5 -quotient of a smooth quintic 3-fold in $\mathbb{P}^4$ obtained as follows. Consider the action of the cyclic group $\mu_5 = \langle \zeta \rangle$ by $(1, \zeta, \zeta^2, \zeta^3, \zeta^4)$ on $\mathbb{P}^4$ and take $\mathbb{P}_{\Delta_1} := \mathbb{P}^4/\mu_5$. The Gorenstein toric Fano variety $\mathbb{P}_{\Delta_1}$ contains exactly 5 isolated singular points. A non-degenerate hypersurface $\overline{W}_1 \subset \mathbb{P}_{\Delta_1}$ is smooth, i.e. $\overline{W}_1 = \widehat{W}_1$, it has the Hodge numbers $h^{1,1}(\overline{W}_1) = 1$, $h^{2,1}(\overline{W}_1) = 21$, and $\pi_1(\overline{W}_1) \cong \mathbb{Z}/5\mathbb{Z}$. This is the single example in the case $|N/N'_{\Delta_i^*}| = 5$.

Similarly one obtains a free μ_3 -quotient $\overline{W}_2$ of a smooth hypersurface of bidegree $(3, 3)$ in $\mathbb{P}^2 \times \mathbb{P}^2$ using the action of the cyclic group $\mu_3 = \langle \zeta \rangle$ on $\mathbb{P}^2 \times \mathbb{P}^2$ by $(1, \zeta, \zeta^2)$ on each $\mathbb{P}^2$. The toric variety $\mathbb{P}_{\Delta_2} := \mathbb{P}^2 \times \mathbb{P}^2/\mu_3$ has 9 isolated singularities. A non-degenerate hypersurface $\overline{W}_2 = \widehat{W}_2 \subset \mathbb{P}_{\Delta_2}$ is smooth, it has the Hodge numbers $h^{1,1}(\overline{W}_2) = 2$, $h^{2,1}(\overline{W}_2) = 29$, and $\pi_1(\overline{W}_2) \cong \mathbb{Z}/3\mathbb{Z}$.

We remark that the dual reflexive polytopes $\Delta_1^*, \dots, \Delta_{16}^*$ have at most 8 vertices. Δ_{14}^* is the single dual reflexive polytope with 8 vertices. The corresponding Gorenstein toric Fano variety $\mathbb{P}_{\Delta_{14}}$ is $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1/\mu_2$ where the cyclic group $\mu_2 = \langle \zeta \rangle$ acts by $(1, \zeta)$ on each $\mathbb{P}^1$.

We collected the information about all 16 reflexive polytopes $\Delta_1, \dots, \Delta_{16}$ in the table below. In this table $\mathcal{P}_\Delta = |\Delta \cap M|$, $\mathcal{V}_\Delta = |\text{vert}(\Delta)|$, $\mathcal{P}_{\Delta^*} = |\Delta^* \cap N|$, and $\mathcal{V}_{\Delta^*} = |\text{vert}(\Delta^*)|$ denote numbers of points and vertices, respectively. Since any reflexive polytope $\Delta_i \subset M_\mathbb{R}$ is uniquely determined by its dual polytope $\Delta_i^* \subset N_\mathbb{R}$, we prefer to use the small number of vertices $v_1, \dots, v_{\mathcal{V}_{\Delta^*}}$ in Δ_i^* in order to describe Δ_i^* by $\mathcal{V}_{\Delta^*} - 4$ independent linear relations among them.

The lattice N is generated by the vertices $v_1, \dots, v_{\mathcal{V}_{\Delta^*}}$ and by an additional vector v expressed as a rational linear combination of the vertices $\{v_i\}$. The sublattice $N'_{\Delta_i^*}$ is generated by $v_1, \dots, v_{\mathcal{V}_{\Delta^*}}$ except for 4 cases: $i = 4, 9, 15, 16$. In the latter cases $N'_{\Delta_i^*}$ is generated by $v_1, \dots, v_{\mathcal{V}_{\Delta^*}}$ and by the vector $2v$.

n°	$ \pi_1 $	$\mathcal{P}_\Delta \mathcal{V}_\Delta$	$\mathcal{P}_{\Delta^*} \mathcal{V}_{\Delta^*}$	$h^{1,1} \ h^{2,1}$	$\chi(\widehat{W})$	vertices $\{v_i\}$ of Δ^* , $N := \mathbb{Z}v + \sum_i \mathbb{Z}v_i$
1	5	26 5	6 5	1 21	-40	$v_1 + v_2 + v_3 + v_4 + v_5 = 0$ $v = \frac{1}{5}(v_2 + 2v_3 + 3v_4 + 4v_5)$
2	3	34 9	7 6	2 29	-54	$v_1 + v_2 + v_3 = v_4 + v_5 + v_6 = 0$ $v = \frac{1}{3}(v_2 + 2v_3 + v_5 + 2v_6)$
3	3	49 5	7 5	2 38	-72	$3v_1 + 3v_2 + v_3 + v_4 + v_5 = 0$ $v = \frac{1}{3}(v_1 + 2v_2 + v_3 + 2v_4)$
4	2	53 5	9 5	3 43	-80	$4v_1 + v_2 + v_3 + v_4 + v_5 = 0$ $v = \frac{1}{4}(2v_1 + v_2 + 2v_3 + 3v_4)$
5	2	77 7	9 6	3 59	-112	$4v_1 + 2v_2 + v_3 + v_4 = 0$ $2v_1 + v_5 + v_6 = 0$ $v = \frac{1}{2}(v_1 + v_2 + v_3 + v_5)$
6	2	77 9	9 7	3 59	-112	$2v_1 + v_2 + v_3 = 2v_1 + v_4 + v_5 = 0$ $2v_1 + v_6 + v_7 = 0$ $v = \frac{1}{2}(v_1 + v_2 + v_4 + v_6)$
7	2	101 5	9 5	3 75	-144	$8v_1 + 4v_2 + 2v_3 + v_4 + v_5 = 0$ $v = \frac{1}{2}(v_1 + v_2 + v_3 + v_4)$
8	2	101 6	9 6	3 75	-144	$4v_1 + 2v_2 + v_3 + v_4 = 0$ $4v_1 + 2v_2 + v_5 + v_6 = 0$ $v = \frac{1}{2}(v_1 + v_2 + v_3 + v_5)$
9	2	29 8	9 6	4 28	-48	$v_1 + v_2 + v_3 + v_4 = v_5 + v_6 = 0$ $v = \frac{1}{4}(v_2 + 2v_3 + 3v_4 + 2v_6)$
10	2	53 8	9 6	4 44	-80	$4v_1 + 2v_2 + v_3 + v_4 = v_5 + v_6 = 0$ $v = \frac{1}{2}(v_1 + v_2 + v_3 + v_5)$
11	2	53 10	9 7	4 44	-80	$2v_1 + v_2 + v_3 = 0$ $2v_1 + v_4 + v_5 = 0$ $v_6 + v_7 = 0$ $v = \frac{1}{2}(v_1 + v_2 + v_4 + v_6)$
12	2	41 9	9 6	4 36	-64	$2v_1 + v_2 + v_3 = 0$ $2v_4 + v_5 + v_6 = 0$ $v = \frac{1}{2}(v_1 + v_2 + v_4 + v_5)$
13	2	41 12	9 7	4 36	-64	$2v_1 + v_2 + v_3 = 0$ $v_4 + v_5 = v_6 + v_7 = 0$ $v = \frac{1}{2}(v_1 + v_2 + v_4 + v_6)$
14	2	41 16	9 8	4 36	-64	$v_1 + v_2 = v_3 + v_4 = 0$ $v_5 + v_6 = v_7 + v_8 = 0$ $v = \frac{1}{2}(v_1 + v_3 + v_5 + v_7)$
15	2	29 5	9 5	5 29	-48	$2v_1 + 2v_2 + 2v_3 + v_4 + v_5 = 0$ $v = \frac{1}{4}(v_1 + 2v_2 + 3v_3 + 2v_5)$
16	2	29 6	9 6	5 29	-48	$v_1 + v_2 + v_3 + v_4 = 0$ $v_3 + v_4 + v_5 + v_6 = 0$ $v = \frac{1}{4}(v_1 + 3v_2 + 2v_4 + 2v_6)$

3. THE BRAUER GROUP OF TORIC HYPERSURFACES

Let X be a smooth quasi-projective variety over $\mathbb{C}$. The cohomological Brauer group $B'(X)$ of X is defined as the torsion subgroup in $H_{\text{et}}^2(X, \mathcal{O}_X^*)$, where $\mathcal{O}_X^*$ denotes the sheaf of units in $\mathcal{O}_X$. If $B(X)$ is the Brauer group of sheaves of Azumaya algebras on X , then there exist a canonical injective homomorphism $B(X) \rightarrow B'(X)$. Recently Kresch and Vistoli proved that this homomorphism is in fact an isomorphism [16].

Let us summarize some well-known results about the Brauer group.

Theorem 3.1. *The subgroup $B(X)_r \subset B(X)$ of elements $x \in B(X)$ with $rx = 0$ can be included into the short exact sequence*

$$0 \rightarrow \text{Pic}(X)/r\text{Pic}(X) \rightarrow H_{\text{et}}^2(X, \mu_r) \rightarrow B(X)_r \rightarrow 0,$$

where μ_r denotes the subsheaf on r -th roots of unity in $\mathcal{O}_X^*$. More generally, there is a short exact sequence

$$0 \rightarrow \text{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{Q}/\mathbb{Z} \rightarrow H_{\text{et}}^2(X, \mathbb{Q}/\mathbb{Z}) \rightarrow B(X) \rightarrow 0,$$

where $H_{\text{et}}^2(X, \mathbb{Q}/\mathbb{Z})$ coincides with the singular cohomology group $H^2(X, \mathbb{Q}/\mathbb{Z})$ in the usual topology of X as a manifold over $\mathbb{C}$.

Proof. See [13] II, Theorem 3.1. □

Corollary 3.2. *If X is a smooth projective variety such that $\text{Pic}(X) \cong H^2(X, \mathbb{Z})$, then one has the isomorphism*

$$B(X) \cong \text{Tors}(H^3(X, \mathbb{Z})).$$

In particular, the last statement holds for smooth Calabi-Yau d -folds ($d \geq 3$).

Proof. Follows from the universal coefficients theorem. □

Theorem 3.3. *Let X be a smooth quasi-projective variety and $Y \subset X$ a closed subvariety. Then the natural homomorphism $B(X) \rightarrow B(X \setminus Y)$ is always injective. This homomorphism is bijective if Y has codimension ≥ 2 . Moreover, if Y is a union of closed irreducible subvarieties $Y_1, \dots, Y_n$ of codimension 1, then one has the exact sequence*

$$0 \rightarrow B(X) \rightarrow B(X \setminus Y) \rightarrow \bigoplus_{i=1}^n H_{\text{et}}^1(Y_i, \mathbb{Q}/\mathbb{Z}),$$

where $H_{\text{et}}^1(Y_i, \mathbb{Q}/\mathbb{Z})$ coincides with the singular cohomology group $H^1(Y_i, \mathbb{Q}/\mathbb{Z})$ in the usual topology if Y_i is smooth.

Proof. See [13] III, Corollary 6.2. □

The Brauer group of a d -dimensional algebraic torus $\mathbb{T}^d$ over $\mathbb{C}$ was computed e.g. by Magid in [21]:

Theorem 3.4. *Using the canonical isomorphism between the cohomology ring $H^*(\mathbb{T}^d, \mathbb{Z})$ and the exterior algebra $\Lambda^* M$ of the $\mathbb{Z}$ -module M , one has the following canonical isomorphisms*

$$B(\mathbb{T}^d) \cong H^2(\mathbb{T}^d, \mathbb{Q}/\mathbb{Z}) \cong \Lambda^2 M \otimes_{\mathbb{Z}} \mathbb{Q}/\mathbb{Z} \cong \text{Hom}(\Lambda^2 M, \mathbb{Q}/\mathbb{Z}).$$

The elements in the Brauer group $B(\mathbb{T}^d)$ can be explicitly described as follows. Take an integer $r > 1$ and two characters $\alpha, \beta \in M \subset \mathcal{O}_{\mathbb{T}^d}^*$ of the algebraic torus $\mathbb{T}^d$. The **symbol algebra** $(\alpha, \beta)_r$ is generated by two elements x, y with relations

$$x^r = \alpha, y^r = \beta, xy = \zeta yx,$$

where ζ is a primitive r -th root of unity. If $r = 2$, then $(\alpha, \beta)_2$ is the classical quaternionic algebra associated with $\alpha, \beta \in \mathcal{O}_{\mathbb{T}^d}^*$.

Let $B(\mathbb{T}^d)_r$ be r -torsion elements in $B(\mathbb{T}^d)$, then $B(\mathbb{T}^d)_r$ is a free $\mathbb{Z}/r\mathbb{Z}$ module with the basis

$$(m_i, m_j)_r, \quad 1 \leq i < j \leq d,$$

where $m_1, \dots, m_d$ is a $\mathbb{Z}$ -basis of the lattice M , i.e.

$$B(\mathbb{T}^d)_r = \Lambda^2(M) \otimes \mathbb{Z}/r\mathbb{Z}.$$

Theorem 3.5. *Let $\Delta \subset M_{\mathbb{R}}$ be a d -dimensional lattice polytope and let $W \subset \mathbb{T}^d$ be a nondegenerate affine hypersurface defined by the equation*

$$f(t) = \sum_{m \in \Delta \cap M} c_m t^m = 0.$$

Then for $d \geq 4$ the natural homomorphism

$$j^* : B(\mathbb{T}^d) \rightarrow B(W)$$

defined by the embedding $j : W \hookrightarrow \mathbb{T}^d$ is an isomorphism.

Proof. By 3.1, we obtain the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Pic}(\mathbb{T}^d) \otimes_{\mathbb{Z}} \mathbb{Q}/\mathbb{Z} & \longrightarrow & H^2(\mathbb{T}^d, \mathbb{Q}/\mathbb{Z}) & \longrightarrow & B(\mathbb{T}^d) \longrightarrow 0 \\ & & \downarrow j_P^* & & \downarrow j_H^* & & \downarrow j^* \\ 0 & \longrightarrow & \text{Pic}(W) \otimes_{\mathbb{Z}} \mathbb{Q}/\mathbb{Z} & \longrightarrow & H^2(W, \mathbb{Q}/\mathbb{Z}) & \longrightarrow & B(W) \longrightarrow 0 \end{array}$$

By 1.3, j_H^* is an isomorphism. Moreover, $\text{Pic}(\mathbb{T}^d) = 0$ since $\mathbb{T}^d$ is a Zariski open subset of $\mathbb{C}^d$. Therefore it suffices to prove that $\text{Pic}(W) \otimes_{\mathbb{Z}} \mathbb{Q} = 0$ (since by surjectivity of $\text{Pic}(W) \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow \text{Pic}(W) \otimes_{\mathbb{Z}} \mathbb{Q}/\mathbb{Z}$ would imply that $\text{Pic}(W) \otimes_{\mathbb{Z}} \mathbb{Q}/\mathbb{Z} = 0$). Using toric resolution of singularities, one can compactify W to a smooth projective semiample hypersurface $\widehat{W}$ in some smooth projective toric variety $\widehat{\mathbb{P}}$ such that $Z := \widehat{W} \setminus W$ is a union of smooth irreducible divisors. By the vanishing theorem for semiample line bundles on toric varieties, we obtain that

$$h^1(\widehat{W}, \mathcal{O}_{\widehat{W}}) = \dots = h^{d-2}(\widehat{W}, \mathcal{O}_{\widehat{W}}) = 0.$$

In particular, $\text{Pic}(\widehat{W}) \cong H^2(\widehat{W}, \mathbb{Z})$. Therefore,

$$\dim_{\mathbb{Q}} \text{Pic}(W) \otimes_{\mathbb{Z}} \mathbb{Q} = \dim_{\mathbb{C}} H^2(\widehat{W}, \mathbb{C}) = b_2(\widehat{W}) = b_{2d-4}(\widehat{W}).$$

Let $Z_1, \dots, Z_k$ be irreducible components of $Z = \widehat{W} \setminus W$. Using the exact sequence

$$\bigoplus_{i=1}^k \mathbb{Q}[Z_i] \rightarrow \text{Pic}(\widehat{W}) \otimes \mathbb{Q} \rightarrow \text{Pic}(W) \otimes \mathbb{Q} \rightarrow 0,$$

it remains to show that the classes $[Z_1], \dots, [Z_k] \in \text{Pic}(\widehat{W})$ generate $\text{Pic}(\widehat{W}) \otimes \mathbb{Q}$. The latter follows from the Poincaré duality together with the long exact sequence for cohomology with compact supports

$$\dots \rightarrow H_c^{2d-4}(W, \mathbb{Q}) \rightarrow H_c^{2d-4}(\widehat{W}, \mathbb{Q}) \rightarrow H_c^{2d-4}(Z, \mathbb{Q}) \rightarrow \dots,$$

where $H_c^{2d-4}(Z, \mathbb{Q}) \cong \mathbb{Q}^k$, $H_c^{2d-4}(\widehat{W}, \mathbb{Q}) \cong \text{Hom}(\text{Pic}(\widehat{W}), \mathbb{Q})$, because the Hodge component of the Hodge type $(d-2, d-2)$ in $H_c^{2d-4}(W, \mathbb{Q})$ is trivial (see [10]).

□

Let us recall the computation of the Brauer group of a smooth (quasi-projective) toric variety $\mathbb{P}_{\Sigma}$ obtained by Demeyer and Ford [11].

By 3.3, $B(\mathbb{P}_{\Sigma}) \cong B(\mathbb{P}_{\Sigma^{(1)}})$ where $\Sigma^{(1)}$ is the subfan of all cones of dimension ≤ 1 in Σ . So we can assume without loss of generality that $\Sigma = \Sigma^{(1)}$ and $\mathbb{P}_{\Sigma}$ is a toric compactification of $\mathbb{T}^d$ by divisors $D_1^{\circ}, \dots, D_n^{\circ}$ ($D_i^{\circ} \cong \mathbb{T}^{d-1}$) corresponding to lattice vectors $e_1, \dots, e_n \in N$. By 3.3, we have the exact sequence

$$0 \rightarrow B(\mathbb{P}_{\Sigma}) \rightarrow B(\mathbb{T}^d) \rightarrow \bigoplus_{i=1}^n H^1(D_i^{\circ}, \mathbb{Q}/\mathbb{Z}).$$

For any $i = 1, \dots, n$ we have the canonical isomorphisms

$$H^1(D_i^{\circ}, \mathbb{Q}/\mathbb{Z}) \cong \text{Hom}(N/\mathbb{Z}e_i, \mathbb{Q}/\mathbb{Z}) \cong M_{e_i} \otimes \mathbb{Q}/\mathbb{Z},$$

where

$$M_{e_i} := \{m \in M : \langle x, e_i \rangle = 0\}.$$

Therefore, one obtains

$$B(\mathbb{T}^d) \supset B(\mathbb{P}_{\Sigma}) = \bigcap_{i=1}^n \text{Ker } \varphi_i,$$

where φ_i denotes the ramification map

$$\varphi_i : B(\mathbb{T}^d) \rightarrow H^1(D_i^{\circ}, \mathbb{Q}/\mathbb{Z})$$

which associates to a given symbol algebra $(\alpha, \beta)_r$ its ramification along the divisor $D_i^{\circ} \subset \mathbb{P}_{\Sigma}$, i.e. a cyclic Galois covering of D_i° of degree r .

Theorem 3.6. *Using the canonical isomorphisms*

$$B(\mathbb{T}^d) \cong \text{Hom}(\Lambda^2 N, \mathbb{Q}/\mathbb{Z}), \quad H^1(D_i^\circ, \mathbb{Q}/\mathbb{Z}) \cong \text{Hom}(N/\mathbb{Z}e_i, \mathbb{Q}/\mathbb{Z}),$$

one obtains the ramification map φ_i from the homomorphism

$$\psi_i : N/\mathbb{Z}e_i \rightarrow \Lambda^2 N$$

$$y \mapsto y \wedge e_i, \quad \forall y \in N$$

by applying the functor $\text{Hom}(, \mathbb{Q}/\mathbb{Z})$. In particular, the image of ψ_i is the sublattice*

$$\text{Im } \psi_i = N \wedge e_i \subset \Lambda^2 N.$$

Proof. These statements are contained in [11] Lemmas 1.5-1.7. $\square$

Let us define the sublattice $N' := \sum_{i=1}^n \mathbb{Z}e_i \subset N$ and write

$$N/N' \cong \mathbb{Z}/c_1\mathbb{Z} \oplus \cdots \oplus \mathbb{Z}/c_d\mathbb{Z}$$

where $c_1, \dots, c_d$ non-negative integers such that $c_1|c_2|\cdots|c_d$. We can choose a $\mathbb{Z}$ -basis $y_1, \dots, y_d$ of N such that $\{c_i y_i\}_{c_i \neq 0}$ is a $\mathbb{Z}$ -basis of N' . Consider another sublattice

$$N \wedge N' = \sum_{i=1}^n N \wedge e_i = \sum_{i=1}^n \text{Im } \psi_i \subset \Lambda^2 N.$$

Then the elements $\{c_i y_i \wedge y_j\}_{c_i \neq 0, i < j}$ form a $\mathbb{Z}$ -basis of $N \wedge N'$. Therefore,

$$\Lambda^2 N / (N \wedge N') \cong \bigoplus_{i < j} \mathbb{Z}/c_i\mathbb{Z}.$$

Thus, one obtains (see [11], Theorem 1.1):

Theorem 3.7. *In the above notations, the Brauer group of a nonsingular toric variety $\mathbb{P}_\Sigma$ is isomorphic to*

$$\bigoplus_{i < j} \text{Hom}(\mathbb{Z}/c_i\mathbb{Z}, \mathbb{Q}/\mathbb{Z}).$$

In particular, if $N' \subset N$ is a sublattice of finite index, then $\mathbb{P}_\Sigma$ has a finite Brauer group

$$B(\mathbb{P}_\Sigma) \cong \text{Hom}(\Lambda^2 N / (N \wedge N'), \mathbb{Q}/\mathbb{Z}) \cong \bigoplus_{i < j} (\mathbb{Z}/c_i\mathbb{Z})^*.$$

Let $\Delta \subset M_\mathbb{R}$ be a d -dimensional lattice polytope and let $W \subset \mathbb{T}^d$ be a nondegenerate affine hypersurface defined by the equation

$$f(t) = \sum_{m \in \Delta \cap M} c_m t^m = 0.$$

Denote by $\overline{W}$ the closure of W in the projective toric variety $\mathbb{P}_\Delta$ and by $\widehat{W}$ a smooth projective desingularization of $\overline{W}$ defined by the fan $\Sigma \subset N_\mathbb{R}$ which is a subdivision of the normal fan to Δ . We denote by $N_\Delta^{(2)}$ the sublattice in N generated by all lattice points $v \in N$ such that the minimum of the

linear function $\langle *, v \rangle$ on Δ is attained on a face $\theta \subset \Delta$ of dimension ≥ 2 . Our main result in this section is the following:

Theorem 3.8. *In the above notation, for $d \geq 4$ the Brauer group $B(\widehat{W})$ of the toric hypersurface $\widehat{W}$ is a cyclic group isomorphic to*

$$\mathrm{Hom}(\Lambda^2 N / (N \wedge N_{\Delta}^{(2)}), \mathbb{Q}/\mathbb{Z}).$$

Proof. Let $e_1, \dots, e_n$ be primitive lattice generators of 1-dimensional cones in Σ such that the minimum of the linear function $\langle *, v \rangle$ on Δ is attained on a face $\theta \subset \Delta$ of dimension ≥ 1 . Denote by $\Sigma' \subset \Sigma^{(1)}$ the subfan of Σ consisting of 0 and all 1-dimensional cones $\mathbb{R}_{\geq 0}e_1, \dots, \mathbb{R}_{\geq 0}e_n$. We set $\widehat{W}^{(1)} = \widehat{W} \cap \mathbb{P}_{\Sigma'}$. Since the complement to $\widehat{W}^{(1)}$ in $\widehat{W}$ has codimension ≥ 2 , by 3.3. it suffices to compute the Brauer group $B(\widehat{W}^{(1)})$. By 3.3, we have the commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & B(\mathbb{P}_{\Sigma'}) & \longrightarrow & B(\mathbb{T}^d) & \longrightarrow & \bigoplus_{i=1}^n \varphi_i H^1(D_i^\circ, \mathbb{Q}/\mathbb{Z}) \\ & & \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow \\ 0 & \longrightarrow & B(\widehat{W}^{(1)}) & \longrightarrow & B(W) & \longrightarrow & \bigoplus_{i=1}^n \phi_i H^1(Z_i^\circ, \mathbb{Q}/\mathbb{Z}) \end{array}$$

By 3.5, β is an isomorphism. In order to understand the homomorphisms α and γ we divide the set of lattice vectors $\mathcal{E} = \{e_1, \dots, e_n\}$ into a union of two disjoint subsets $\mathcal{E}_1 \cup \mathcal{E}_2$. The set $\mathcal{E}_2$ consists of all $e_i \in \mathcal{E}$ such that the minimum of $\langle *, e_i \rangle$ on Δ is attained on a face $\theta \subset \Delta$ of dimension ≥ 2 and we define $\mathcal{E}_1 := \mathcal{E} \setminus \mathcal{E}_2$.

For any $e_i \in \mathcal{E}$ we denote by θ_i the maximal subface of Δ on which the minimum of $\langle *, e_i \rangle$ on Δ is attained.

If $e_i \in \mathcal{E}_2$, then $d_i := \dim \theta_i \geq 2$ and the divisor Z_i° is isomorphic to the product $V_i \times \mathbb{T}^{d-d_i-1}$, where V_i is a θ_i -nondegenerate affine hypersurface in a torus $\mathbb{T}^{d_i}$. By 1.3, the ramification homomorphism

$$\phi_i : B(W) \rightarrow H^1(Z_i^\circ, \mathbb{Q}/\mathbb{Z})$$

can be identified with the composition of the ramification homomorphism

$$\varphi_i : B(\mathbb{T}^d) \rightarrow H^1(D_i^\circ, \mathbb{Q}/\mathbb{Z})$$

with the monomorphism

$$H^1(D_i^\circ, \mathbb{Q}/\mathbb{Z}) \rightarrow H^1(Z_i^\circ, \mathbb{Q}/\mathbb{Z}).$$

Thus, we obtain

$$B := \bigcap_{e_i \in \mathcal{E}_2} \mathrm{Ker} \varphi_i = \bigcap_{e_i \in \mathcal{E}_2} \mathrm{Ker} \phi_i.$$

Using 3.6, we can describe the dual group B^* as cokernel of the homomorphisms

$$\psi_i : N/\mathbb{Z}e_i \rightarrow \Lambda^2 N, \quad e_i \in \mathcal{E}_2,$$

i.e. we have

$$\sum_{e_i \in \mathcal{E}_2} \text{Im } \psi_i = N \wedge N_{\Delta}^{(2)} \text{ and } B^* = N / (N \wedge N_{\Delta}^{(2)}).$$

Therefore, it remains to prove that

$$B = \bigcap_{e_i \in \mathcal{E}} \text{Ker } \phi_i = B(\widehat{W}^{(1)}),$$

i.e., that for any $e_i \in \mathcal{E}_1$ the image $\text{Im } \phi_i = \text{Im } \gamma \circ \psi_i$ is contained in $N \wedge N_{\Delta}^{(2)}$. Now we consider two cases for $e_i \in \mathcal{E}_1$:

Case 1: Let $e_i \in N_{\Delta}^{(2)}$. Then the image of ϕ_i is contained in $N \wedge e_i \subset N \wedge N_{\Delta}^{(2)}$.

Case 2: Let $e_i \notin N_{\Delta}^{(2)}$. Then the image of ϕ_i is a sublattice $N_i \wedge e_i$, where N_i is a direct summand of N of rank $d-1$ containing e_i (N_i is a sublattice of rank $d-1$ which is orthogonal to the 1-dimensional face θ_i). The smoothness of $\widehat{W}$ implies that both lattices N_i and $N_{\Delta}^{(2)}$ contain a direct summand of N of rank $d-2$. Therefore, $N_i/N_i \cap N_{\Delta}^{(2)}$ is a cyclic group and we have

$$N_i = \mathbb{Z}v + N_i \cap N_{\Delta}^{(2)}$$

for some lattice vector $v \in N_i$ which is proportional to e_i . By $v \wedge e_i = 0$, we have

$$N_i \wedge e_i \subset (\mathbb{Z}v + N_i \cap N_{\Delta}^{(2)}) \wedge e_i \subset N_{\Delta}^{(2)} \wedge e_i \subset N \wedge N_{\Delta}^{(2)}.$$

Thus, all divisors Z_i ($e_i \in \mathcal{E}_1$) do not define non-trivial ramification conditions on the the subgroup B of the Brauer group $B(W)$ and

$$B \cong B(\widehat{W}^{(1)}) \cong B(\widehat{W}) \cong \text{Hom}(\Lambda^2 N / (N \wedge N_{\Delta}^{(2)}), \mathbb{Q}/\mathbb{Z}).$$

□

Corollary 3.9. *Let $\widehat{W}$ be a smooth projective Calabi-Yau d -fold obtained as a crepant desingularization of a nondegenerate projective hypersurface $\overline{W} \subset \mathbb{P}_{\Delta}$ corresponding to a reflexive polytope Δ of dimension $d \geq 4$. Then the Brauer group $B(\widehat{W})$ is a cyclic group isomorphic to*

$$\text{Hom}(\Lambda^2 N / (N \wedge N_{\Delta^*}''), \mathbb{Q}/\mathbb{Z}),$$

where $N_{\Delta^*}'' \subset N$ is the sublattice generated by all lattice points in $N \cap \Delta^*$ which are contained in faces of Δ^* of codimension > 2 .

Proof. The statement follows immediately from 3.8 and from the easy combinatorial equality $N_{\Delta}^{(2)} = N_{\Delta^*}''$ using the duality between faces of dual reflexive polytopes $\Delta \subset M_{\mathbb{R}}$ and $\Delta^* \subset N_{\mathbb{R}}$. □

4. THE BRAUER GROUP OF CALABI-YAU 3-FOLDS

Let Δ be a 4-dimensional reflexive polytope. Then

$$N''_{\Delta^*} := \sum_{v \in \theta^* \cap N, \dim \theta^* = 1} \mathbb{Z}v \subset N$$

is a sublattice of finite index. We write

$$N/N''_{\Delta^*} = \mathbb{Z}/c_1\mathbb{Z} \oplus \mathbb{Z}/c_2\mathbb{Z} \oplus \mathbb{Z}/c_3\mathbb{Z} \oplus \mathbb{Z}/c_4\mathbb{Z},$$

where $c_1|c_2|c_3|c_4$. In this case $c_1 = c_2 = 1$ and

$$B(\widehat{W}) \cong \text{Hom}(\Lambda^2 N / (N \wedge N''_{\Delta^*}), \mathbb{Q}/\mathbb{Z}) \cong \mathbb{Z}/c_3\mathbb{Z}.$$

Using the Calabi-Yau data [18], one can check all 473 800 776 reflexive polytopes Δ and find exactly 16 reflexive polytopes of dimension 4 with $c_3 > 1$. For each of these 16 polytopes, one obtains

$$N/N''_{\Delta^*} \cong \mathbb{Z}/p\mathbb{Z} \oplus \mathbb{Z}/p\mathbb{Z}, \quad p = 2, 3, 5.$$

So we have again 16 families with the cyclic Brauer group $\mathbb{Z}/p\mathbb{Z}$. The main observation is that these 16 families are exactly the mirrors of 16 non-simply connected families of Calabi-Yau 3-folds with the cyclic fundamental group $\mathbb{Z}/p\mathbb{Z}$ ($c = 2, 3, 5$) (see Section 2). Therefore, we can exchange the lattices $N \leftrightarrow M$ and polytopes $\Delta^* \leftrightarrow \Delta$ so that for all 4-dimensional reflexive polytopes we have the isomorphisms

$$(2) \quad \Lambda^2 M / (M \wedge M''_{\Delta}) \cong N / N'_{\Delta^*} \quad \text{and} \quad \Lambda^2 N / (N \wedge N''_{\Delta^*}) \cong M / M'_{\Delta}$$

Unfortunately we do not have any natural combinatorial explanation of the isomorphisms (2).

The sublattice $N''_{\Delta^*} \subset N$ defines a (ramified) Galois covering $\widehat{W}'' \rightarrow \widehat{W}$ with the Galois group $G = N/N''_{\Delta^*} \cong \mathbb{Z}/p\mathbb{Z} \oplus \mathbb{Z}/p\mathbb{Z}$. If we consider G as a 2-dimensional vector space over the finite field $\mathbb{F}_p$, then the Brauer group $B(\widehat{W})$ is isomorphic to $\text{Hom}(\Lambda^2 G, \mathbb{Q}/\mathbb{Z})$, i.e. to a 1-dimensional $\mathbb{F}_p$ -space. This allows to look at $B(\widehat{W})$ as the discrete torsion group $H^2(G, U(1))$ for the orbifold $\widehat{W}''/G$.

We finish this section by describing a typical example $\widehat{W}_1^*$ of a Calabi-Yau 3-fold from our list (the mirror of $\widehat{W}_1$ from Section 2) with Brauer group $B(\widehat{W}_1^*) \cong \mathbb{Z}/5\mathbb{Z}$. This example gives a 1-parameter family of Calabi-Yau 3-folds $\widehat{W}_1^*$ having a close connection to the counterexample to the global Torelli considered by Aspinwall-Morrison and B. Szendrői [1, 27].

Consider the action of the group $G := \mu_5 \times \mu_5$ on $\mathbb{P}^4$ by

$$\begin{aligned} \zeta_1 &\rightarrow (1, \zeta_1, \zeta_1^2, \zeta_1^3, \zeta_1^4) \\ \zeta_2 &\rightarrow (1, \zeta_2, \zeta_2^3, \zeta_2, 1). \end{aligned}$$

This action is not free and the quotient $\mathbb{P}_{\Delta_1^*} := \mathbb{P}^4/G$ has cyclic singularities of type $\frac{1}{5}(3, 1, 1)$ along 10 curves. Let $\widehat{W}_1^*$ be the crepant desingularization of a generic Calabi-Yau hypersurface $\overline{W}_1^* \subset \mathbb{P}_{\Delta_1^*}$. Then

$$\begin{aligned}\pi_1(\widehat{W}_1^*) &= 0 = B(\widehat{W}_1) \\ B(\widehat{W}_1^*) &= \Lambda^2 G \cong \mu_5 \cong \pi_1(\widehat{W}_1).\end{aligned}$$

REFERENCES

- [1] P. S. Aspinwall, D. R. Morrison, *Chiral Rings Do Not Suffice: $N=(2,2)$ Theories with Nonzero Fundamental Group*, Phys.Lett. B334 (1994), 79-86.
- [2] P. S. Aspinwall, D. R. Morrison (with an appendix by M. Gross), *Stable Singularities in String Theory*, Commun.Math.Phys. 178 (1996), 115-134.
- [3] V. V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*. J. Algebraic Geom. 3 (1994), no. 3, 493-535.
- [4] V. V. Batyrev, *Stringy Hodge numbers of varieties with Gorenstein canonical singularities*. Integrable systems and algebraic geometry (Kobe/Kyoto, 1997), 1-32, World Sci. Publishing, River Edge, NJ, 1998.
- [5] V. V. Batyrev, L.A. Borisov, *Mirror duality and string-theoretic Hodge numbers*. Invent. Math. 126 (1996), no. 1, 183-203.
- [6] L. A. Borisov, L. Chen, G.G. Smith, *The orbifold Chow ring of toric Deligne-Mumford stacks*. J. Amer. Math. Soc. 18 (2005), no. 1, 193-215.
- [7] I. Brunner, J. Distler, *Torsion D-Branes in Nongeometrical Phases* Adv.Theor.Math.Phys. 5 (2002), 265-309.
- [8] I. Brunner, J. Distler, R. Mahajan, *Return of the Torsion D-Branes*, Adv.Theor.Math.Phys. 5 (2002), 311-352.
- [9] V.I. Danilov, *The geometry of toric varieties*, Uspekhi Mat. Nauk 33 (1978), no. 2(200), 85-134, 247.
- [10] V.I. Danilov, A.G. Khovanskii, *Newton polyhedra and an algorithm for calculating Hodge-Deligne numbers*. Izv. Akad. Nauk SSSR Ser. Mat. 50 (1986), no. 5, 925-945.
- [11] F. Demeyer, T. Ford *On the Brauer group of toric varieties*. Trans. Amer. Math. Soc. 335 (1993), 559-577.
- [12] M. Gross, *Special Lagrangian fibrations. I*. Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds (Cambridge, MA, 1999), 65-93, Amer. Math. Soc., Providence, RI, 2001.
- [13] A. Grothendieck *Le groupe de Brauer I,II,III*. In: J. Giraud (ed) et al.: Dix exposés sur la cohomologie des schémas, pp. 46-189. North-Holland, Amsterdam (1968).
- [14] C. Haase, B. Nill, *Lattices generated by skeletons of reflexive polytopes*, in preparation.
- [15] A. Klemm, M. Kreuzer, E. Riegler, E. Scheidegge, *Topological String Amplitudes, Complete Intersection Calabi-Yau Spaces and Threshold Corrections*, hep-th/0410018.
- [16] Andrew Kresch, Angelo Vistoli, *On coverings of Deligne-Mumford stacks and surjectivity of the Brauer map*, math.AG/0301249.
- [17] M. Kreuzer, H. Skarke, *Complete classification of reflexive polyhedra in four dimensions*, Adv.Theor.Math.Phys. 4 (2002) 1209-1230, hep-th/0002240.
- [18] M. Kreuzer, H. Skarke, <http://hep.itp.tuwien.ac.at/~kreuzer/CY.html>
- [19] M. Kreuzer, H. Skarke, *PALP: A package for analyzing lattice polytopes with applications to toric geometry*, Computer Physics Commun. 157 (2004) 87-106, math.NA/0204356.

- [20] B. Nill, *Complete toric varieties with reductive automorphism group*, math.AG/0407491.
- [21] A. R. Magid, *Brauer groups of linear algebraic groups with characters*. Proc. Amer. Math. Soc. 71 (1978), no. 2, 164–168.
- [22] M. Oka, *On the topology of the Newton boundary II*, J. Math. Soc. Japan, Vol **32**, n.1 (1980), 65–92.
- [23] M. Oka, *A Lefschetz type theorem in a torus*. Singularity theory (Trieste, 1991), 574–593, World Sci. Publishing, River Edge, NJ, 1995.
- [24] M. Oka, *Non-degenerate complete intersection singularity*. Actualits Mathematiques. Hermann, Paris, 1997.
- [25] T. Pantev, E. Sharpe, *String compactifications on Calabi-Yau stacks*, hep-th/0502044.
- [26] T. Pantev, E. Sharpe, *GLSM's for gerbes (and other toric stacks)*, hep-th/0502053.
- [27] B. Szendrői, *On an example of Aspinwall and Morrison*, Proc. AMS, vol 132, n. 3 (2003), 621–632.
- [28] C. Vafa, E. Witten *Orbifolds with discrete torsion*, J. Geom. Phys. **15** (1995)

MATHEMATISCHES INSTITUT, UNIVERSITÄT TÜBINGEN, AUF DER MORGENSTELLE 10,
72076 TÜBINGEN, GERMANY

E-mail address: `victor.batyrev@uni-tuebingen.de`

INSTITUT FÜR THEORETISCHE PHYSIK, TECHNISCHE UNIVERSITÄT WIEN, WIEDNER
HAUPTSTR. 8-10/136, A-1040 VIENNA, AUSTRIA

E-mail address: `maximilian.kreuzer@tuwien.ac.at`